

Model of the aral sea sedimentary cycles

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© SGEM2018. The goal of this project was to create a detailed model of the Aral Sea sedimentary cycles. Modern methods of image processing and wavelet analysis were used to reveal the nature of lamination and determine the average and interval sedimentation rates. The results obtained were then compared with those from the radiocarbon dating. The images used in this project are photographs of core (sediment) samples taken with hollow plastic tubes. The samples were removed from the tubes by cutting them in half lengthwise, and then photographed. The image resolution was 0.34 mm. The task of identifying the periodic components and determining the quantitative characteristics of individual cycles and microcycles was accomplished through the wavelet analysis of the “brightness” curve. The cycles that make up the sedimentation model have periods of 1, ≈ 2 , ≈ 5 , ≈ 13 , ≈ 50 and ≈ 200 years (i.e. they are well aligned with the common climatic cycles). The average sedimentation rate is estimated at 2-3 mm/year. The model shows that a 50 cm thick sediment forms in ~ 160 years. The radiocarbon dating, however, produced a different outcome – 440 ± 100 years. Obviously, the differences in the results are not trivial even if the likely level of error is taken into account.

<http://dx.doi.org/10.5593/sgem2018/1.1/S01.037>

Keywords

Annual laminations, Depositional cycle, Image processing, Thin-laminated clays, Wavelet

References

- [1] Deevey, E.S., M.S., Gross, G.E., Hutchinson, and H.L. Kraybill, The natural ^{14}C contents of materials from hard-water lakes, *Proc. Nat. Acad. Sci.*, 40, 285-288, 1954.
- [2] Bronk Ramsey, C., Radiocarbon calibration and analysis of stratigraphy: The OxCal program, *Radiocarbon*, 37(2), 425-430, 1995.
- [3] Botvinkina L.N. Methodological Guide for the Study of Laminations. M.: “Nauka”, 1965, p. 259.
- [4] Korn H. Schichtung und absolute Zeit.-Rundschau, 1935, 26.
- [5] Boomer, I., N. Aladin, I. Plotnikov, and R. Whatley, The paleolimnology of the Aral Sea: a review. *Quarter. Sci. Rev.*, 19, 1259-1278, 2000.
- [6] Tarasov, P.E., S.P. Harrison, L. Saarse, et al., Lake Status Records from the FSU//Database Documentation Version 2. IGBP PAGES/World Data Center-A for Paleoclim. Ser. 96-032. 1996. www.ftp.ngdc.noaa.gov/paleo/aleolimnology/lakelevels/former_ussr
- [7] [7] Brodskaya N.G. Bottom sediments and sedimentation processes of the Aral Sea. M.: Academy of Sciences of the USSR, 1952, p. 103.
- [8] Khrustalyov Yu.P., Reznikov S.A., Turovskiy D.S. Lithology and geochemistry of the Aral Sea sediments. — Rostov-on-Don, 1977, p. 159.

- [9] Astaf'eva N. M. Wavelet analysis: basic theory and some applications. Phys. Usp. 39 1085–1108 (1996); DOI: 10.1070/PU1996v039n11ABEH000177
- [10] Nourgaliev, D. K., Khasanov, D. I. (1992). Solar cycles recorded in Late Permian sediments. Solnechnye dannye 8, pp. 82-85.